

Optically Well-Studied Clusters  
With Cool Gas Temperatures

GRANT NAG5-2157

Final Report

For the period 15 January 1993 through 14 January 1994

Principal Investigator  
Dr. John P. Huchra

September 1994

Prepared for  
**National** Aeronautics and Space Administration  
Greenbelt, Maryland 20771

Smithsonian Institution  
Astrophysical Observatory  
Cambridge, Massachusetts 02138

The Smithsonian Astrophysical Observatory  
is a member of the  
Harvard-Smithsonian Center for Astrophysics

N95-70181

Unclass

29/89 0025978

(NASA-CR-196937) OPTICALLY  
WELL-STUDIED CLUSTERS WITH COOL GAS  
TEMPERATURES Final Report, 15 Jan.  
1993 - 14 Jan. 1994 (Smithsonian  
Astrophysical Observatory) 4 p

11N-89-12  
0017  
25978  
p 4



We obtained and analyzed a 3000 second PSPC observation of the southern cluster of galaxies S753. Although the flat-field, exposure corrected image gave only a hint of diffuse X-ray emission from an intracluster medium (see Figure 1), smoothing this region on larger scales showed the presence and extent of a hot ICM (see Figure 2). We used the smoothed image to define sources and regions for spectral analysis.

We excluded point sources and measured the X-ray spectrum of the ICM in four annular regions, to an outer radius of 16'. We compared the spectra to Raymond models and found that the gas is isothermal (within the uncertainties) with a temperature of about 2 keV, with possibly a cooler component in the inner 2' (from a possible cooling flow). We also measured the X-ray surface brightness profile. Under the reasonable assumption that the gas is in hydrostatic equilibrium, we used the gas density profile derived from the surface brightness profile and an ICM temperature of 2.5 keV to derive the total cluster mass as a function of cluster radius. From optical observations, we have obtained velocities for about twelve galaxies in this poor cluster. We will use these to make a second estimate of the total cluster mass for comparison with our X-ray measurements.

A paper on this short analysis is in preparation for submission to the *PASP*. Proposal funding was used entirely to support the visit of L. DaCosta to CfA.

s753\_r36 - S753  
(IRAF)




Figure 1. Photon map of our 3000 second PSPC exposure of S753.

s753\_sm2 - S753  
(IRAF)

306.5 319.0 0.1005





Figure 2. Smoothed intensity profile from Figure 1.

